CHAPTER 8: CONFIDENCE INTERVAL ESTIMATES for Means and Proportions

Introduction: We want to know the value of a parameter for a population. We don't know the value of this parameter for the entire population because we don't have data for the entire population. (If we did already know the value of the parameter, we wouldn't need to do any statistical investigation or calculations.) We will use sample statistics to estimate population parameters.

Recall from chapter 2:	A parameter is	
------------------------	----------------	--

If we don't know the value of a population parameter, we can estimate it using a sample statistic.

Recall from chapter 2: A statistic is______

Using data from a sample to draw a conclusion about a population is called ______ statistics.

Two Types of Estimates for population parameters:

1) **POINT ESTIMATE: A population parameter can be estimated by one number: the sample statistic**. This is called a **point estimate**.

(Statistical theory has identified desirable properties of point estimates, which are studied in more depth in upper level statistics classes. One property usually considered desirable is that a point estimate be "unbiased", meaning that the average of the point estimates from all possible samples would equal the true value of the population parameter.)

The "best" point estimate of a population mean μ is

The "best" point estimate of a population proportion p is______

The "best" point estimate of a population standard deviation σ is______

2) CONFIDENCE INTERVAL ESTIMATE:

- The population parameter is estimated by an **interval of numbers** that we believe contains the true (unknown) value of the population parameter.
- We are able to state how confident we are that the interval estimate contains the true (unknown) value of the parameter.
- This **confidence interval estimate** is built using two items: a point estimate, and margins of error; the margins of error are also called error bounds.

We will use confidence interval estimates based on sample data to estimate

- a population average (mean)
- population proportion

Confidence intervals for means and proportions are symmetric; the point estimate is at the center of the interval. The endpoints of the interval are found as

(point estimate – error bound , point estimate + error bound)

(For some other parameters, such as standard deviation, a confidence interval may not be symmetric about the point estimate, moving different distances above and below the point estimate to the ends of the interval estimate.)

We'll learn by example to calculate the point estimates and the error bounds and what they mean. The last 3 pages of these notes has a concise summary of formulas, procedures, and interpretations. We'll start in class by examining a jar with beads to determine the proportion of beads in the jar that are blue; after we explore the concepts, then we'll move on to the mathematical calculations.

CHAPTER 8CONFIDENCE INTERVAL ESTIMATE for an
unknown POPULATION PROPORTION p

a. Statistics and data in this example are based on information from : http://sf.streetsblog.org/2014/08/15/car-free-households-are-booming-in-san-francisco/ http://en.wikipedia.org/wiki/List of U.S. cities with most households without a car

A trend in urban development is to reduce the need for residents to have a car; city neighborhoods are often ranked for "walkability". In recent studies, the US city with the lowest car ownership rate is New York City; a majority (56%) of households are "car-free" with only 44% of households owning any vehicles. San Jose has the highest car ownership rate of large US cities; only about 6% of households "car-free". San Francisco's percent of "car-free" households has changed rapidly in recent years.

Suppose a recent study of 1200 households in San Francisco showed that 372 households were "car-free". Construct and interpret a 95% confidence interval for the true proportion of households in San Francisco that are "car-free". Use a 95% confidence level.

population parameter: p = _____

random variable p' = _____

We are using sample data to estimate an unknown proportion for the whole population

HOW TO CALCULATE THE CONFIDENCE INTERVAL			
Point Estimate = p '	Confidence Level CL is area in the middle		
Error Bound = (Critical Value)(Standard Error) $\mathbf{EBP} = Z_{\alpha/2} \sqrt{\frac{\mathbf{p'q'}}{n}}$	Critical Value is $Z_{\alpha/2}$ is the Z value that creates area of CL in the middle; Z ~N(0,1)	Standard Error	
Confidence Interval = Point Estimate + Error Bound Confidence Interval = $p' \pm EBP$	Use POSITIVE value of Z invnorm (<i>area to left</i> , 0, 1)	$\sqrt{\frac{p q}{n}}$	

Calculations and interpretation in context of the problem:

CHAPTER 8 EXAMPLE 2: Using Confidence Intervals

Before elections, many polls survey samples of likely voters to try to gauge support for the candidates for public office in the election. Since we can't poll all voters before the election, the polls are using the sample data to try to predict the proportion of the population of all actual voters who will vote for each candidate.

2A. Suppose a poll indicates 52% of those in survey indicating they intend to vote for Candidate A for President. The poll uses a 95% confidence level and has a margin of error of $\pm 4\%$. Construct the confidence interval estimate for candidate A. Based on this confidence interval, does the poll give an indication (with 95% confidence) whether Candidate A will have more than 50% of the vote.

Interval:

Explanation of Results:

2B. Suppose a different poll indicates 55% of those in survey indicating they intend to vote for Candidate A for President. The poll uses a 95% confidence level and has a margin of error of $\pm 3\%$. Construct the confidence interval estimate for candidate A. Based on this confidence interval, does the poll give an indication (with 95% confidence) whether Candidate A will have more than 50% of the vote.

Interval:

Explanation of Results:

2C. In Example 1 the point estimate is _____ = _____; the confidence interval is ______

(i) Can we conclude with 95% confidence that more than 25% of SF households are "car-free"?

(ii) Can we conclude with 95% confidence that more than 30% of SF households are "car-free"?

2D. What does it mean when we say the confidence level is 95% or that "we are 95% confident"?

CHAPTER 8CONFIDENCE INTERVAL ESTIMATE for unknown POPULATION MEAN μEXAMPLE 3:when the POPULATION STANDARD DEVIATION σ is KNOWN

A soda bottling plant fills cans labeled to contain 12 ounces of soda. The filling machine varies and does not fill each can with exactly 12 ounces. To determine if the filling machine needs adjustment, each day the quality control manager measures the amount of soda per can for a random sample of 50 cans. Experience shows that its filling machines have a known population standard deviation of 0.35 ounces.

In today's sample of 50 cans of soda, the sample average amount of soda per can is 12.1 ounces.

- a. Construct and interpret a 90% confidence interval estimate for the true population average amount of soda contained in all cans filled today at this bottling plant. Use a 90% confidence level.
- X = _____

population parameter: $\mu =$ _____

random variable $\overline{\mathbf{X}}$ = ______

We are using sample data to estimate an unknown mean (average) for the whole population

HOW TO CALCULATE THE CONFIDENCE INTERVAL for µ			
When σ IS known, use the Standard normal distribution Z ~ N(0,1)			
Point Estimate = $\overline{\mathbf{X}}$	Confidence Level CL is area in the middle	Standard	
Error Bound = (Critical Value)(Standard Error)	Critical Value is Z_{α} is the Z value that	Error	
$\mathbf{EBM} = Z_{\alpha/2} \left(\frac{\sigma}{\sqrt{n}}\right)$	creates area of CL in the middle; $Z \sim N(0,1)$	σ	
Confidence Interval = Point Estimate + Error Bound	Use POSITIVE value of Z		
Confidence Interval = $\overline{X} \pm EBM$	$\operatorname{invnorm}(area to left, 0, 1)$	\sqrt{n}	

Calculations and interpretation in context of the problem:

CHAPTER 8CONFIDENCE INTERVAL ESTIMATE for unknown POPULATION MEAN μ EXAMPLE 3:when the POPULATION STANDARD DEVIATION σ is NOT KNOWN

a. The traffic commissioner wants to know the average speed of all vehicles driving on River Rd. Police use radar to observe the speeds for a sample of 20 vehicles on River Rd.

For the vehicles in the sample, the average speed is 31.3 miles per hour with standard deviation 7.0 mph.

Construct and interpret a 98% confidence interval estimate of the true population average speed of all vehicles on River Rd. Use a 98% confidence level.

X = _____

population parameter: $\mu =$ _____

random variable $\overline{\mathbf{X}}$ =

We are using sample data to estimate an unknown mean (average) for the whole population

HOW TO CALCULATE THE CONFIDENCE INTERVAL for µ			
When σ is NOT known, use the <i>t</i> distribution with degrees of freedom = sample size -1 : <i>t</i> with df = n -1)			
Point Estimate = $\overline{\mathbf{X}}$	Confidence Level CL is area in the middle		
Error Bound = (Critical Value)(Standard Error)	Critical Value $t_{\alpha/}$ is the <i>t</i> value that	Standard	
$\mathbf{EBM} = \mathbf{t}_{\alpha/2} \left(\frac{\mathbf{s}}{\sqrt{n}} \right)$	/2 creates an area of CL in the middle;	Error	
Confidence Interval = Point Estimate + Error Bound Confidence Interval = $\overline{X} + EBM$	Use POSITIVE value of t TI-84: $t_{\alpha/2} = invT(area to left, df)$	$\frac{3}{\sqrt{n}}$	

Calculations and interpretation in context of the problem:

b. In Example 3, suppose that you were not given the sample mean and sample standard deviation and instead you were given a list of data for the speeds (in miles per hour) of the 20 vehicles.
19 19 22 24 25 27 28 37 35 30 37 36 39 40 43 30 31 36 33 35 How would you use the data to do this problem?

NOTE: Use of t-distribution requires the underlying population of individual values to be approximately normally distributed. It is OK if this assumption is violated a little, but if the underlying population of individual values has a distribution that differs too much from the normal distribution, then this confidence method is not appropriate, and statisticians would use other techniques that we do not study in Math 10.

CHAPTER 8 Working Backwards: Finding the Error Bound and Point Estimate EXAMPLE 4: if we know only the confidence interval:

Jessie wants to estimate the average cost of a ride to the airport, from her apartment, using Uber and using Lyft and using a taxi.

For Uber, the 90% confidence interval estimate of the average fare is \$20 to \$27.

For Lyft, the 90% confidence interval estimate of the average fare is \$21 to \$25

For a taxi, the 90% confidence interval estimate of the average fare is \$28 to \$31

(NOTE: This information is based on random samples of fare estimates generated by fare comparison websites for Uber, Lyft, and taxis using one particular address and the closest major airport to that address, with requests made at a variety of times. This may not be representative of the relationships between average fares for Lyft and Uber and Taxi in all locations.)

a. Find the point estimate for the true average fare for each. Explain how to use the confidence interval to find the point estimate. Which has the lowest point estimate?

b. Find the margin of error (error bound for each). Explain how to use the confidence interval to find the error bound. Which has the lowest margin of error?

c. Can we draw any conclusions about the true average fare based on the estimates above?

CHAPTER 8: Confidence Interval for a Proportion: Calculating the Sample Size needed in a Study

Given a desired confidence level and a desired margin of error, how large a sample is needed?

$$\text{EBP} = Z_{\alpha/2} \sqrt{\frac{p'q'}{n}}$$

We know the error bound EBP that we want.

We know the confidence level CL we want, so we can find Z_{α_2} corresponding to the desired CL.

We don't know p' or q' until we do the study, so we will assume for now that $p' = q' = \frac{1}{2} = 0.5$

Then we can substitute all these values into EBP = $Z_{\alpha/2} \sqrt{\frac{p'q'}{n}}$ and solve for n.

Solving EBP =
$$Z_{\alpha/2} \sqrt{\frac{p'q'}{n}}$$
 for n gives $n = \left(\frac{Z_{\alpha/2}}{EBP}\right)^2 p'q$

Sample Size Formula	$(\mathbf{z})^2$	The 0.25 appears in the formula because
to determine sample size n	$\mathbf{n} = \left(\frac{Z_{\alpha/2}}{2} \right) (.25)$	we are assuming that $p' = q' = \frac{1}{2} = 0.5$
needed when estimating a		AT WAVE POUND UP to the payt higher integer
population proportion p		ALWAIS KOUND OF to the next higher integer

EXAMPLE 5: Finding the Sample Size:

a. Public opinion and political polls often do surveys with a 95% confidence level and 3% margin of error. Find the sample size needed.

$$\mathbf{n} = \left(\frac{Z_{\alpha/2}}{EBP}\right)^2 (.25) =$$

b. Suppose a margin of error of 2% was wanted with a 95% confidence level. Find the sample size needed.

$$\mathbf{n} = \left(\frac{Z_{\alpha/2}}{EBP}\right)^2 (.25) =$$

c. Suppose a margin of error of 3% was wanted with a 90% confidence level. Find the sample size needed.

$$\mathbf{n} = \left(\frac{Z_{\alpha/2}}{EBP}\right)^2 (.25) =$$

d. Suppose a poll uses a sample size of n=100, and a confidence level of 95%. Estimate the expected error bound using $p' = q' = \frac{1}{2} = 0.5$

 $\text{EBP} = Z_{\alpha/2} \sqrt{\frac{\mathbf{p'q'}}{n}}$

Note the actual error bound will differ after the study is done because we will know p' and q' and will no longer be estimating that p'=q'=0.5

e. Is the error bound in part d large or small compared to the examples in parts a, b, c? Explain why this happened.

CHAPTER 8 EXTRA PRACTICE EXAMPLES : CALCULATING CONFIDENCE INTERVALS

Sources for Practice Examples: #6 based on Chapter 8 Practice in Introductory Statistics from OpenStax available for download for free at <u>http://cnx.org/content 11562/latest/</u>.; #7based on information from http://alloveralbany.com/archive/2008/06/10/how-big-is-a-scoop-of-ice-cream; #8 based on information from http://www.pewsocialtrends.org/2016/05/24/for-first-time-in-modern-era-living-with-parents-edges-out-other-living-arrangements-for-18-to-34-year-olds/; #9 based on information from http://www.pewinternet.org/2016/02/11/15-percent-of-american-adults-have-used-online-dating-sites-or-mobile-dating-apps/

PRACTICE EXAMPLE 6: A supermarket chain is deciding what produce providers to purchase from. A sample of 20 heads of lettuce is selected to estimate the average weight of the lettuce from this provider. The population standard deviation for the weight is known to be 0.2 pounds. The sample of 20 heads of lettuce had a mean weight of 2.2 pounds with a sample standard deviation of 0.1 pounds.

Calculate and interpret a confidence interval estimate for the true average weight of all heads of lettuce from this provider. Use a 90% confidence level.

PRACTICE EXAMPLE 7: The staff at the All Over Albany (N.Y.) website wanted to know how much ice cream is in "a scoop". For a random sample of 5 ice cream stores, the amount of ice cream contained in a scoop was 6.8, 3.9, 5.4, 3.55, 5.2 ounces.

Calculate and interpret a 95% confidence interval estimate of the average amount of ice cream in a scoop for all ice cream sold at ice cream stores in Albany NY. Use a 95% confidence level.

PRACTICE EXAMPLE 8: Suppose a survey of 300 young adults age 18 to 24 showed that 81 had used online dating. Calculate and interpret a confidence interval estimate for the true proportion of all young adults age 18 to 24 who ever used online dating. Use a 95% confidence level.

PRACTICE EXAMPLE 9: Suppose a survey of 500 people age 18 to 34 indicated that 32.2% of them live with one or both of their parents. Calculate and interpret a confidence interval estimate for the true proportion of all people age 18 to 34 who live with one or both parents. Use a 94% confidence level.

CHAPTER 8: FLOW CHART VIEW OF FORMULAS FOR CONFIDENCE INTERVAL ESTIMATES © 2015 R Bloom For more details, see the presentation of the formulas in boxes for each case on the next page.





CHAPTER 8: CONFIDENCE INTERVALS: SUMMARY OF FORMULAS, PROCEDURES, & INTERPRETATIONS

Confidence Interval for a Proportion p To estimate a population proportion p (binomial probability of success) Point Estimate + Margin of Error (Margin of Error is also called Error Bound Point Estimate: Sample Proportion: $p' = \frac{x}{n} = \frac{number \ of \ successes \ in \ sample}{total \ number \ in \ sample}$ Error Bound: EBP = (critical value)(standard error) = $Z_{\alpha/2} \sqrt{\frac{p'q'}{2}}$ The critical value Z_{α_2} depends on the confidence level. Z_{α_2} is the Z value that puts an area equal to the confidence level (CL) in the middle of standard normal distribution N(0,1) $Z_{\alpha/2}$ tells us how many "appropriate standard deviations" to enclose about the point estimate, where the "standard error" $\sqrt{\frac{p'q'}{n}}$ is the appropriate standard deviation for a proportion $p' \pm EBP$ which is $p' \pm Z_{\alpha/2} \sqrt{\frac{p'q'}{n}}$ **Confidence Interval:**

Confidence Interval for a Mean μ when σ is known

To estimate the population average μ if we already know the population standard deviation σ .

Point Estimate + Margin of Error (Margin of Error is also called Error Bound)

Point Estimate: Sample Average (Sample Mean) \overline{X}

Error Bound: EBM = (critical value)(standard error) = $Z_{\alpha/2} \left(\frac{\sigma}{\sqrt{n}}\right)$

The critical value $Z_{\alpha/2}$ depends on the confidence level. $Z_{\alpha/2}$ is the Z value that puts an area equal to the

confidence level (CL) in the middle of standard normal distribution N(0,1)

 Z_{α_2} tells us how many "appropriate standard deviations" to enclose about the point estimate, where the "standard error" $\frac{\sigma}{\sqrt{n}}$ is the appropriate standard deviation for the sample mean

-	
1	

 $\overline{\mathbf{X}} \pm Z_{\alpha/2} \left(\frac{\sigma}{\sqrt{n}} \right)$ \overline{X} + EBM Confidence Interval: which is

Confidence Interval for a Mean μ when σ is NOT known

To estimate the population average μ and we do not know the population standard deviation σ . Use the sample standard deviation s to estimate the population standard deviation σ

Point Estimate + Margin of Error (Margin of Error is also called Error Bound)

Point Estimate: Sample Average (Sample Mean) \overline{X}

Error Bound: EBM = (critical value)(standard error) = $t_{\alpha/2} \left(\frac{s}{\sqrt{n}}\right)$

The critical value $t_{\alpha_{\alpha}}$ depends on the confidence level. $t_{\alpha_{\alpha}}$ is the t value that puts an area equal to the

confidence level (CL) in the middle of the student t -distribution with n - 1 degrees of freedom

 $t_{\alpha/2}$ tells us how many "appropriate standard deviations" we need to move away from the point estimate, where $\frac{s}{\sqrt{n}}$ is an *estimate* of the standard error ("appropriate standard deviation") for the sample mean

 $\overline{\mathbf{X}} \pm \text{EBM}$ which is $\overline{\mathbf{X}} \pm \mathbf{t}_{\alpha/2} \left(\frac{\mathbf{s}}{\sqrt{n}}\right)$ Confidence Interval:

YOU ARE NOT PERMITTED TO BRING A PRINTOUT OF THIS PAGE AS YOUR NOTES FOR AN EXAM OR QUIZ. You CAN write whatever information you want from this page into your handwritten notes for exams or guizzes. © 2016 R. Bloom

CHAPTER 8: CONFIDENCE INTERVALS: SUMMARY OF FORMULAS, PROCEDURES, & INTERPRETATIONS

Interpreting the Confidence Interval for a PROPORTION (2 ways to word it)

We estimate with _____% confidence that the true <u>proportion</u> of the population that *describe the population parameter in the situation of this problem* is between _____ and _____

We estimate with _____% confidence that between _____% and _____% of the population *describe the population parameter in the situation of this problem*

Interpreting the Confidence Interval for a MEAN (average)

We estimate with _____% confidence that the true population <u>average</u> (or mean) *describe the population parameter in the situation of this problem* is between ______ and _____

What is the meaning of the Confidence Level? What does it mean to be CL% confident?

The confidence level represents an "expected accuracy rate" for the confidence interval process. It tells us what percent of confidence intervals calculated in this manner would be "good estimates".

If we took repeated samples and calculated many confidence interval estimates (one for each sample), we expect that CL% of the confidence interval estimates would be "good estimates" that successfully enclose (capture) the true value of the population parameter that we want to estimate.

If we took repeated samples), we expect that 100% - CL% of the confidence interval estimates would be "bad estimates" that would NOT enclose (capture) the true value of the population parameter.

Note that the confidence interval does not estimate individual data values. It estimates proportions or averages. It does NOT imply that CL% of the data lies within the confidence interval.

Finding the Point Estimate and Error Bound (Margin of Error) if we know the Confidence Interval:The interval is (lower bound, upper bound)Point Estimate = (lower bound + upper bound)/2Error Bound = (upper bound - lower bound)/2



CL specifies the area in the middle

 $\alpha = 1 - CL$ is area "outside", split equally between both tails

 $\frac{\alpha}{2}$ is area in one tail. df = degrees of freedom = n-1

To find
$$t_{\alpha/2}$$
: TI-84+: invT $(1-\frac{\alpha}{2}, df)$

OR use invT ($\frac{\alpha}{2}$, df) and take absolute value (drop the "-" sign)



TI-83,83+: Use INVT program; ask instructor to download it to your calculator: PRGM INVT enter area to the left and df after prompts: area to left is $1-\frac{\alpha}{2}$; (if using $\frac{\alpha}{2}$ as area to left, drop the "-" sign)

Without calculator or if calculator does not have inverse t: Use a student's-t distribution probability table. Value of t is found at the intersection of the column for the confidence level and row for degrees of freedom

YOU ARE <u>NOT</u> PERMITTED TO BRING A <u>PRINTOUT</u> OF THIS PAGE AS YOUR NOTES FOR AN EXAM OR QUIZ. You CAN write whatever information you want from this page into your handwritten notes for exams or quizzes. © 2016 R. Bloom